

# DEXAMOLCOLD® NIGHT CARE

Patient package insert according to Pharmacists' Regulations (Preparations) 1986

Carefully read the entire package insert before using this medication.

Composition: Each caplet contains: Paracetamol 500 mg, Pseudoephedrine hydrochloride 25 mg, Dextromethorphan hydrobromide 10 mg, Chlorpheniramine maleate 2 mg

List of inactive ingredients: Povidone, croscarmellose sodium (approx. 1 mg sodium), magnesium stearate, silica colloidal anhydrous, hypromellose, titanium dioxide, macrogol, carnauba wax pigments: E-127, E-132, E-104, E-133

Therapeutic action: Dexamolcold night care contains four active ingredients: Paracetamol is an analgesic and an antipyretic. Pseudoephedrine hydrochloride is a sympathomimetic amine which relieves runny nose, nasal and ear congestion, accompanying cold and/or sinusitis. Dextromethorphan hydrobromide suppresses cough accompanying cold. Chlorpheniramine maleate is an antihistamine that dries mucus.

Indications: This medicine is intended for the symptomatic relief of cold, cough and nasal congestion accompanied by fever and pain. Night care medicine

When should this medicine not be used?

Do not use this medicine if you are pregnant or breastfeeding. Do not use this medicine if you are aware of any hypersensitivity to one of its ingredients. Do not use this medicine if you are taking monoamine oxidase inhibitors (MAOI) (for the treatment of depression, for instance) or within 14 days of stopping treatment with MAOIs. Do not use this medicine for the relief of chronic cough.

Do not use this medicine without consulting your physician before commencing treatment: if you suffer, or have suffered in the past, from impaired function of the respiratory system (e.g. asthma, cough that occurs with too much phlegm, chronic bronchitis), the heart and/or vascular system, hypertension, coronary arterial disease, increased intraocular pressure (glaucoma), the liver, the kidney/urinary tract, alcoholism, jaundice, the thyroid, the nervous system, enlargement of the prostate gland, trouble urinating, diabetes.

How will this medicine influence your daily schedule?

Use of this medicine may impair alertness. It is therefore advisable to take special care when driving, operating dangerous machinery or when engaging in any other activity which requires alertness. Children should be cautioned against bicycle riding or playing near traffic, etc.

Do not drink wine or other alcoholic beverages during the period of treatment with this medicine.

Warnings: Paracetamol may cause damage in the liver in the following cases: when administered at a dosage higher than the recommended one, when administered for extended periods, when consuming alcoholic beverages during the treatment period, when taking additional medicines that affect the function of the liver. Do not use this medicine frequently without consulting your physician (also see below - "Dosage"). If you are aware of any sensitivities to foods or drugs inform your physician before taking this medicine.

Patients above the age of 60 years should notify their physician before taking this medicine as they might be sensitive to decongestive drugs.

Avoid exceeding the recommended dosage while fasting. Do not take additional medicines for reduction of fever and relief of pain, or other cold medicines, without consulting your physician or pharmacist in order to avoid paracetamol overdose/toxicity.

Drug interactions: If you are taking other medicines concurrently, or have just completed treatment with another medicine inform your physician in order to prevent hazards or inefficacy arising from drug interactions, especially in regard to: drugs affecting the central nervous system (e.g. appetite suppressants, tranquilizers, sedatives and hypnotics, for Parkinson's disease, epilepsy, anti-allergics, surgical anaesthetics and narcotic analgesics); anticoagulants; antidepressants; antihypertensives and drugs for the heart; drugs that increase liver enzyme

See next page

# דקסמולקולד® לטיפול בלילה

נشرة طبية للمستهلك بموجب انظمة الصيدلة (مستحضرات) ١٩٨٦

الرائي النشرة يامعان حتى نهايتها قبل استعمالك للدواء

Composition: Each caplet contains: Paracetamol 500 mg, Pseudoephedrine hydrochloride 25 mg, Dextromethorphan hydrobromide 10 mg, Chlorpheniramine maleate 2 mg

List of inactive ingredients: Povidone, croscarmellose sodium (approx. 1 mg sodium), magnesium stearate, silica colloidal anhydrous, hypromellose, titanium dioxide, macrogol, carnauba wax pigments: E-127, E-132, E-104, E-133

Therapeutic action: Dexamolcold night care contains four active ingredients: Paracetamol is an analgesic and an antipyretic. Pseudoephedrine hydrochloride is a sympathomimetic amine which relieves runny nose, nasal and ear congestion, accompanying cold and/or sinusitis. Dextromethorphan hydrobromide suppresses cough accompanying cold. Chlorpheniramine maleate is an antihistamine that dries mucus.

Indications: This medicine is intended for the symptomatic relief of cold, cough and nasal congestion accompanied by fever and pain. Night care medicine

When should this medicine not be used?

Do not use this medicine if you are pregnant or breastfeeding. Do not use this medicine if you are aware of any hypersensitivity to one of its ingredients. Do not use this medicine if you are taking monoamine oxidase inhibitors (MAOI) (for the treatment of depression, for instance) or within 14 days of stopping treatment with MAOIs. Do not use this medicine for the relief of chronic cough.

Do not use this medicine without consulting your physician before commencing treatment: if you suffer, or have suffered in the past, from impaired function of the respiratory system (e.g. asthma, cough that occurs with too much phlegm, chronic bronchitis), the heart and/or vascular system, hypertension, coronary arterial disease, increased intraocular pressure (glaucoma), the liver, the kidney/urinary tract, alcoholism, jaundice, the thyroid, the nervous system, enlargement of the prostate gland, trouble urinating, diabetes.

How will this medicine influence your daily schedule?

Use of this medicine may impair alertness. It is therefore advisable to take special care when driving, operating dangerous machinery or when engaging in any other activity which requires alertness. Children should be cautioned against bicycle riding or playing near traffic, etc.

Do not drink wine or other alcoholic beverages during the period of treatment with this medicine.

Warnings: Paracetamol may cause damage in the liver in the following cases: when administered at a dosage higher than the recommended one, when administered for extended periods, when consuming alcoholic beverages during the treatment period, when taking additional medicines that affect the function of the liver. Do not use this medicine frequently without consulting your physician (also see below - "Dosage"). If you are aware of any sensitivities to foods or drugs inform your physician before taking this medicine.

Patients above the age of 60 years should notify their physician before taking this medicine as they might be sensitive to decongestive drugs.

Avoid exceeding the recommended dosage while fasting. Do not take additional medicines for reduction of fever and relief of pain, or other cold medicines, without consulting your physician or pharmacist in order to avoid paracetamol overdose/toxicity.

Drug interactions: If you are taking other medicines concurrently, or have just completed treatment with another medicine inform your physician in order to prevent hazards or inefficacy arising from drug interactions, especially in regard to: drugs affecting the central nervous system (e.g. appetite suppressants, tranquilizers, sedatives and hypnotics, for Parkinson's disease, epilepsy, anti-allergics, surgical anaesthetics and narcotic analgesics); anticoagulants; antidepressants; antihypertensives and drugs for the heart; drugs that increase liver enzyme

See next page

# דקסמולקולד® לטיפול בלילה

עלון לצרכן לפי תקנות הרוקחים (תכשירים) התשמ"ו - 1986

יש לקרוא בעיון את העלון עד סופו לפני השימוש בתרופה.

הרכב: כל קפליה מכילה:

Composition: Each caplet contains: Paracetamol 500 mg, Pseudoephedrine hydrochloride 25 mg, Dextromethorphan hydrobromide 10 mg, Chlorpheniramine maleate 2 mg

List of inactive ingredients: Povidone, croscarmellose sodium (approx. 1 mg sodium), magnesium stearate, silica colloidal anhydrous, hypromellose, titanium dioxide, macrogol, carnauba wax pigments: E-127, E-132, E-104, E-133

Therapeutic action: Dexamolcold night care contains four active ingredients: Paracetamol is an analgesic and an antipyretic. Pseudoephedrine hydrochloride is a sympathomimetic amine which relieves runny nose, nasal and ear congestion, accompanying cold and/or sinusitis. Dextromethorphan hydrobromide suppresses cough accompanying cold. Chlorpheniramine maleate is an antihistamine that dries mucus.

Indications: This medicine is intended for the symptomatic relief of cold, cough and nasal congestion accompanied by fever and pain. Night care medicine

When should this medicine not be used?

Do not use this medicine if you are pregnant or breastfeeding. Do not use this medicine if you are aware of any hypersensitivity to one of its ingredients. Do not use this medicine if you are taking monoamine oxidase inhibitors (MAOI) (for the treatment of depression, for instance) or within 14 days of stopping treatment with MAOIs. Do not use this medicine for the relief of chronic cough.

Do not use this medicine without consulting your physician before commencing treatment: if you suffer, or have suffered in the past, from impaired function of the respiratory system (e.g. asthma, cough that occurs with too much phlegm, chronic bronchitis), the heart and/or vascular system, hypertension, coronary arterial disease, increased intraocular pressure (glaucoma), the liver, the kidney/urinary tract, alcoholism, jaundice, the thyroid, the nervous system, enlargement of the prostate gland, trouble urinating, diabetes.

How will this medicine influence your daily schedule?

Use of this medicine may impair alertness. It is therefore advisable to take special care when driving, operating dangerous machinery or when engaging in any other activity which requires alertness. Children should be cautioned against bicycle riding or playing near traffic, etc.

Do not drink wine or other alcoholic beverages during the period of treatment with this medicine.

Warnings: Paracetamol may cause damage in the liver in the following cases: when administered at a dosage higher than the recommended one, when administered for extended periods, when consuming alcoholic beverages during the treatment period, when taking additional medicines that affect the function of the liver. Do not use this medicine frequently without consulting your physician (also see below - "Dosage"). If you are aware of any sensitivities to foods or drugs inform your physician before taking this medicine.

Patients above the age of 60 years should notify their physician before taking this medicine as they might be sensitive to decongestive drugs.

Avoid exceeding the recommended dosage while fasting. Do not take additional medicines for reduction of fever and relief of pain, or other cold medicines, without consulting your physician or pharmacist in order to avoid paracetamol overdose/toxicity.

Drug interactions: If you are taking other medicines concurrently, or have just completed treatment with another medicine inform your physician in order to prevent hazards or inefficacy arising from drug interactions, especially in regard to: drugs affecting the central nervous system (e.g. appetite suppressants, tranquilizers, sedatives and hypnotics, for Parkinson's disease, epilepsy, anti-allergics, surgical anaesthetics and narcotic analgesics); anticoagulants; antidepressants; antihypertensives and drugs for the heart; drugs that increase liver enzyme

See next page

המשך מעבר לדף



